

Optimizing meadow management using temperature sums



In the Upper Rhine region, grasslands are an important feed resource for cattle farms. If properly managed, they can contribute to the food and protein self-sufficiency of livestock farms by providing high-quality forage. The use of benchmarks such as temperature sums should facilitate their exploitation by helping farmers to anticipate the benchmark stages of their grasslands.

Temperature sums: a grassland management tool

Good management of grassland, whether by grazing or harvesting, means **exploiting it at the right stage**, i.e. finding a compromise between yield, feed value and animal needs.

The availability of water, nutrients, sunlight and temperature are essential for plant growth. However, at the end of winter, the primary factor limiting grass growth remains **air temperature**, and this factor will condition the appearance of the different physiological stages of the grasses present in permanent grasslands.

The 5 main reference stages for permanent grasslands are based on the phenological stage of the grasses that make them up: vegetation start, 5cm ear of corn, early heading, full heading, flowering. These stages are associated with meadow management guidelines that optimize both forage quantity and quality: putting to grass, dethatching, early mowing, late mowing, etc. For a given grass, reaching these stages requires a certain level of temperature accumulation, also known as "temperature sum". With climate change, this criterion varies considerably from one year to the next, and calendar date references are becoming obsolete. On the other hand, the use of temperature sums has the advantage of taking into account both the climatic context of the year and the location of the meadow.

The 5 main stages are associated with benchmarks for managing grassland to optimise both forage quantity and quality.




Figure 1: The main phenological stages of forage grasses (from Fourrage Mieux 2014)

A variety of permanent grasslands

Work carried out by INRA Toulouse (Cruz et al., 2010) has made it possible to classify different permanent grasslands according to the functional types of the dominant grasses that make them up. This typology is based on a number of functional traits shared by species of the same type: leaf dry matter content, leaf area, leaf life, flowering date, maximum plant height, leaf resistance to breakage. It allows us to define each species according to its habitat preference and use value (ability to fulfill a given function in a forage system).

| Grassland type | A | B | b | C | D |
|-------------------------------|--|--|--|--|---|
| Soil fertility | Fertile environment | Fertile environment | Fertile environment | Low-fertility environment | Low-fertility environment |
| Plant size | Small | Large | | Small | Medium |
| Precocity | Very early | Fairly early | Quite late | Fairly early | Very late |
| usage | Early and frequent grazing | Early mowing and/or late haying | Mowing and summer grazing | Not very suitable for mowing. Good feed value at vegetative stage. | Typical flora of little-used summer pastures or rangelands. Low forage value |
| Representative species | <i>English ryegrass, woolly coot, sweet vernal grass, timothy, meadow foxtail...</i> | <i>orchardgrass, meadow fescue, tall fescue, Kentucky bluegrass, upright bromegrass...</i> | <i>common bentgrass, timothy, creeping couch grass, meadow bluegrass, soft coot...</i> | <i>cretelle, red fescue, sheep's fescue...</i> | <i>brachypode penné, Loudun oats, nard raide, canche des champs, molinie bleue...</i> |

Table 1: Typology of French permanent grasslands (after Cruz et al. 2010)

By definition, a natural grassland is multi-species, and will never support a single group of species. If a functional type represents more than 66% of all grasses, the prairie will be considered of the same type as the dominant functional type.

Ex: For a meadow where type A, B and C grasses represent respectively 25%, 70% and 5% of the grasses, we consider that the meadow is type B.

If the dominant functional type represents less than 66% of the grasses, the grassland will be classified according to its component types as soon as they represent more than 20% of all grasses.

Ex: For a meadow where type B, b and C grasses represent 45%, 40% and 5% of the grasses respectively, the meadow is considered to be type Bb.



Applying this typology to Swiss meadows

Swiss grasslands are grouped according to four intensities of use, from "intensive" to "extensive", resulting in **13 common grassland types** on the northern slopes of the Alps and in the Jura. They are named after the most representative species, which are mainly indicator plants.

- Intensive use: very frequent mowing/grazing with a high level of fertilization
 - ✓ Italian ryegrass meadow
 - ✓ Mow-and-graze meadow with English ryegrass and Kentucky bluegrass
 - ✓ Meadow foxtail meadow
- Semi-intensive use: mowing/grazing
 - ✓ Cocksfoot meadow
 - ✓ Crested dog's-tail grassland
- Low-intensity use
 - ✓ Orchard grass meadow
 - ✓ Yellow oat meadow
 - ✓ Dandelion pasture
 - ✓ Grassland with bentgrass and red fescue
 - ✓ Moist, rich grassland with 2 subtypes: marram grassland and rush grassland
- Extensive use
 - ✓ Brome grass prairie
 - ✓ Spikenard pasture
 - ✓ Litter meadow with 3 sub-types: mullein meadow, Davall's sedge meadow and brown sedge meadow



| Use | Intensive | | | Mi-intensive | | Low intensity | | | | | Extensive | |
|--------------------------------|-------------------------|--|-----------------------|------------------|------------------------------|----------------------|-------------------|-------------------|---|--------------------|---------------------|-------------------|
| Meadow type | Italian ryegrass meadow | English ryegrass and Kentucky bluegrass meadow | Meadow foxtail meadow | Cocksfoot meadow | Crested dog's-tail grassland | Orchard grass meadow | Yellow oat meadow | Dandelion pasture | Grassland with bentgrass and red fescue | Moist, rich meadow | Brome grass prairie | Spike-nard meadow |
| Main valuation | Mowing | Mowing +grazing | Mowing | Mowing | Pasture | Mowing + grazing | Mowing | Pasture | Mowing+grazing/ Grazing | Pasture | Mowing | Grazing |
| Precocity | Early | Very early | Early | Fairly early | Fairly early | Late | Late | Late | Late | Late | Very late | Late |
| Fertility | Strong | Strong | Strong | Intermediary | Intermediary | | Low | Intermediary | Low | Low | Low | Low |
| Productivity | Strong | Strong | Quite strong | Quite strong | Intermediary | Intermediary | Low | Very Low | Very Low | Very low | Very low | Very low |
| French typology correspondance | A | A | A | B | C | B | C | b | C | D | D | D |

Table 2: Typology of Swiss grasslands and correspondance with French permanent grassland types (from the knowledge platform of the Association pour le développement de la Culture Fourragère eADCF)

Management guidelines adapted to the diversity of grasslands

The typology of permanent grasslands, based primarily on species earliness and soil fertility, has enabled us to define the sums of temperatures required to reach the different benchmark stages for each type of grassland, and thus to anticipate the corresponding management practices to be implemented, both for grazing management and for 1st mowing.

| Meadow type | | A or B | b | C |
|------------------|--------------------------------|-----------------------------------|--------------|-------------------------|
| Precocity | | Very early to early | Late | Fairly early |
| Soil fertility | | High | Intermediate | Low |
| Productivity | | Strong | Average | Low |
| Prairie Stadium | Management practices | Temperature sums (in degrees day) | | |
| Vegetation start | Grazing turnout | 250 to 300 | 400 | 400 |
| 5 cm ear | tillering | 500 | 800 | 800 |
| Ear 10 cm | End 1 ^{er} of grazing | 500 to 600 | 1000 | 900 |
| Early heading | Early mowing: silage, wrapping | 700t o 800 | 1200 | Not suitable for mowing |
| Full heading | Early hay | 800 to 1100 | 1500 | |
| Flowering | Late hay | 900 to 1200 | 1600 | |

Table 3: Correspondence between temperature sums, grass physiological stages and best management practices for permanent grasslands type A or B, b and C

How to calculate temperature sums

Cumulative mean daily temperatures are a good estimator of the different phenological stages of plants. However, a number of calculation rules need to be observed to adapt to the specific context of prairie grasses.

1. *The daily average* is based on the raw minimum and maximum values from the local weather station, without any prior correction.
2. *The vegetation zero* for meadows is the base 0°C.
3. *The daily averages are capped* at 18°C, as the growth acceleration phenomenon is considered to stop above this threshold.
4. *The initialization date* for cumulative daily temperatures is February 01.
5. Altitude leads to slower growth due to colder average daily temperatures. The *altitude difference* between the weather station and the plot is taken into account by applying a correction of -0.6°C to the daily average for every 100-meter rise in altitude.

In practice:

- For each day from 01/02 onwards, calculate the average daily temperature $(\text{Minimum temperature} + \text{Maximum temperature})/2$
- If the average daily temperature is $< 0^\circ\text{C}$, use an average daily temperature of 0°C .
Ex: If the average temperature is -5°C , note 0°C
- If the average daily temperature is between 0°C and 18°C , use this value,
Ex: If the average temperature is 13°C , we note 13°C
- If the average daily temperature is $> 18^\circ\text{C}$, use an average daily temperature of 18°C .
Ex: If the average temperature is 21°C , we note 18°C .
- Add the daily average to obtain the sum of temperatures in degrees days since 01/02




Source : Parc Naturel Régional Lorraine



Benchmarks for different weather stations

The following tables show the average dates at which the various milestones have been reached at selected weather stations over the last ten years (2014-2023).



German weather stations

| Weather station | LAHR (156 m) | | | LENZKIRCH-RUHBÜHL (854 m) | | |
|--|--------------|----------|----------|---------------------------|----------|----------|
| | Median | Min. | Max. | Median | Min. | Max. |
| 300 Degree-days (Grassing) | March 22 | March 11 | 9-Apr. | 19-Apr. | 7-Apr. | 27-Apr. |
| 400 Degree-days | 1-Apr. | March 21 | 15-Apr. | May 2nd | 21-Apr. | May 10 |
| 500 Degree-days (End of trimming - thorn at 5 cm) | 13-Apr. | 5-Apr. | 23-Apr. | May 11 | May 1 | May 21st |
| 700 Degree-days (Early mowing-Early heading) | May 1 | 21-Apr. | May 9th | May 28 | May 21st | June 8 |
| 800 Degree-days | May 8 | 28-Apr. | May 16 | June 5 | May 31st | June 14 |
| 1000 Degree-days (Early hay-Early flowering) | May 22 | May 14 | May 31st | June 18 | June 14 | June 27 |
| 1100 Degree-days | May 29 | May 21st | June 6 | June 24 | June 20 | July 3 |
| 1200 Degree-days (Medium hay-Flowering) | June 4 | May 27 | June 12 | 1-july | June 27 | July 10 |

Table 4: Dates for reaching certain milestones in grassland, based on the sum of mean temperatures base 0-18 accumulated since February 01 over the period 2014-2023 (Source: LKV).



Swiss weather stations

| Weather station | AESCH (382 m) | | | COURTEMELON (450 m) | | | FRICK (390 m) | | |
|--|---------------|----------|----------|---------------------|----------|---------|---------------|----------|---------|
| | Median | Min. | Maxi. | Median | Min. | Max. | Median | Min. | Max. |
| 300 Degree-days (Grassing) | March 19 | March 10 | 11-Apr. | March 27 | March 16 | 14-Apr. | March 25 | March 16 | 13-Apr. |
| 400 Degree-days | March 30 | March 19 | 18-Apr. | 9-Apr. | 1-Apr. | 21-Apr. | 5-Apr. | March 7 | 19-Apr. |
| 500 Degree-days (end of trimming at 5 cm) | 11-Apr. | Apr. 4 | 23-Apr. | 21-Apr. | 11-Apr. | 28-Apr. | 18-Apr. | 8-Apr. | 26-Apr. |
| 700 Degree-days (Early mowing-Early heading) | 29-Apr. | 18-Apr. | May 10 | May 9th | 28-Apr. | May 16 | May 6 | 25-Apr. | May 13 |
| 800 Degree-days | May 9th | 24-Apr. | May 19th | May 17 | May 6 | May 25 | May 15 | May 3 | May 22 |
| 1000 Degree-days (Early hay-Early flowering) | May 24 | May 8 | June 1 | May 30 | May 22 | June 8 | May 28 | May 18 | June 5 |
| 1100 Degree-days | May 30 | May 16 | June 7 | June 5 | May 29 | June 14 | June 2 | May 25 | June 11 |
| 1200 Degree-days (Medium hay-Flowering) | June 5 | May 22 | June 13 | June 11 | June 5 | June 19 | June 8 | June 1 | June 17 |

Table 5: Dates for reaching a number of grassland milestones based on the sum of mean temperatures base 0-18 accumulated since February 01 over the period 2014-2023 (Source: Agrométéo- Agroscope)

French weather stations

| Weather station | BERG (300 m) | | | CARSPACH (332 m) | | | MUNSTER (420 m) | | |
|--|--------------|----------|---------|------------------|----------|---------|-----------------|----------|---------|
| | Median | Min. | Maxi. | Median | Min. | Max. | Median | Min. | Max. |
| 300 Degree-days (Grassing) | March 25 | March 18 | 13-Apr. | March 22 | March 13 | 10-Apr. | March 25 | March 16 | 13-Apr. |
| 400 Degree-days | 6-Apr. | March 31 | 20-Apr. | 1-Apr. | March 25 | 18-Apr. | 6-Apr. | March 30 | 20-Apr. |
| 500 Degree-days (End of trimming - thorn at 5 cm) | 18-Apr. | 7-Apr. | 28-Apr. | 15-Apr. | 7-Apr. | 24-Apr. | 19-Apr. | 8-Apr. | 27-Apr. |
| 700 Degree-days (Early mowing-Early heading) | May 6 | 25-Apr. | May 13 | May 2nd | 22-Apr. | May 9th | May 6 | 26-Apr. | May 13 |
| 800 Degree-days | May 13 | May 4 | May 22 | May 11 | 29-Apr. | May 18 | May 15 | May 4 | May 22 |
| 1000 Degree-days (Early hay-Early flowering) | May 28 | May 20 | June 4 | May 25 | May 15 | June 2 | May 29 | May 20 | June 6 |
| 1100 Degree-days | June 3 | May 27 | June 10 | May 31st | May 22 | June 8 | June 4 | May 27 | June 11 |
| 1200 Degree-days (Medium hay-Flowering) | June 9 | June 2 | June 16 | June 5 | May 28 | June 13 | June 10 | June 3 | June 17 |

Table 6: Dates for reaching certain milestones in grassland based on the sum of mean temperatures base 0-18 accumulated since February 01 over the period 2014-2023 (Source: Météo France)

Relationship between temperature sum, yield and forage quality in 1st cycle

As part of the Obs-Herbe network, the Swiss national grassland observatory set up by Agroscope, 23 intensive permanent grasslands were monitored from 2017 to 2019. On the basis of 4 samples taken at regular intervals between the start of sprouting and first mowing each year, it was possible to measure yields as well as nitrogen, fiber and ash contents, enabling energy and nitrogen values to be calculated and linked to the temperature sums recorded during sampling.

By demonstrating the correlation between temperature sums and yields, as well as between temperature sums and feed values, this work has provided the table below determining yields and feed values as a function of temperature sums.

| Degree-days | Yield (Tonne DM/ha) | Protein (g/kg) | Digestibility (%) | Energy : NEL (MJ/kg) | BREAD (g/kg) |
|-------------|---------------------|----------------|-------------------|----------------------|--------------|
| 550 | 2.89 | 165.2 | 78.6 | 6.5 | 110.0 |
| 600 | 3.24 | 155.9 | 77.4 | 6.4 | 103.7 |
| 650 | 3.58 | 146.6 | 76.2 | 6.3 | 97.4 |
| 700 | 3.93 | 137.4 | 75 | 6.2 | 91.2 |
| 750 | 4.27 | 128.1 | 73.8 | 6 | 84.9 |
| 800 | 4.61 | 118.8 | 72.6 | 5.9 | 78.7 |
| 900 | 5.30 | 100.2 | 70.2 | 5.7 | 66.1 |

NEL: net energy milk - DM: dry matter - g/kg: grams per kilogram - MJ/kg: megajoules per kilogram
 PAIN: absorbable proteins in the intestine synthesized from degradable nitrogenous matter

Table 7: Relationship between temperature sum, yield and forage quality based on 23 Swiss meadows monitored over 3 years (Source: Agroscope)



Source :
 Swiss Agricultural Research

Management of subsequent grazing cycles

To improve grassland management, it's important to consider grass growth patterns throughout the entire season, not only in spring.

This means that, after initial exploitation (severe grazing, mowing), plants are left with only a reduced leaf surface and must therefore mobilize their reserves to produce new leaves. During this period, growth slows considerably. As this leaf surface regenerates, the plants will capture more and more light to produce biomass through photosynthesis. As a result, the yield of the grassland becomes higher and higher, and faster and faster, until the grass reaches the 3-leaf stage. Beyond the 3-leaf stage, grassland growth slows down to zero. In fact, a grass blade generally bears only 3 (or 4) leaves: the appearance of an additional leaf leads to the senescence and death of the oldest leaf.




Figure 2: Grass growth curve (after André Voisin 1957)

The return to previously used grassland (grazing, mowing) should therefore take place neither too early nor too late. Harvesting too early limits the replenishment of plant reserves, leading to dwarfing and eventual disappearance. Conversely, harvesting too late leads to wastage, as the oldest leaves disappear.

Meadow resting time should therefore be based on the 3-leaf stage, and will depend on the season, soil and climate conditions, and the type of meadow. It will be around 20 days in spring, and 35 to 40 days, or even longer depending on water deficit and temperature, in summer and autumn.



Leaf life, a grazing management indicator for the following cycles

In the summer, temperature sums alone are no longer sufficient to control grazing, as very often the factor limiting regrowth is the level of rainfall. However, there is a direct link between grass leaf life and temperature sums. The warmer the weather, the faster the vegetative cycles.

Leaf life is a feature that reflects the recycling of leaf tissue. Leaf life span (LLS) varies from species to species and conditions mowing and grazing practices. Thus, late use of species with short LOS will result in a high loss of biomass through senescence, whereas early use of species with long LOS will penalize their lifespan through a lack of adaptation to too-frequent defoliation.

| Types of grass | Leaf life (in degree-days) |
|----------------|-------------------------------|
| Type A | 800 |
| Type B | 1000 |
| Type b | 830 |
| Type C | 1100 |
| Type D | 1100 |

Table 9: Leaf life (in degree-days) of different types of grass (after Cruz et al, 2010)

In practice, if an RGA leaf dies after 800 degree-days, mowing (or grazing) should be carried out within 40 to 45 days of first harvesting. For orchardgrass or tall fescue, whose leaf life is around 1,000 degree-days, this harvesting cycle should take place within 50 to 60 days after first harvesting.

Conclusion

Grassland management requires a great deal of technical skill, and in particular a good knowledge of the phenological stages of the species that make up the grassland, in order to provide sufficient quantities of quality grass (palatability, digestibility, feed value). Temperature sums are a real decision-making tool to help advisors and farmers identify these stages and implement good grassland management practices. They are available, for example, through the weekly publications of the French Grass Growth Monitoring Networks.



Bibliography

ADCF: "Type de prairies dans la production fourragère " <https://www.eagff.ch/fr/connaitre-les-plantes-des-prairies/types-de-prairies>

Carrère P., Pontes L. da S., Andueza D., Louault F., Rosseel D., et al.. (2010) : "Evolution de la valeur nutritive de graminées prairiales au cours de leur cycle de développement", *Fourrages*, 201, 27-35.

Chambres d'agriculture de Lorraine (2018): Fiche technique Pâturage tournant

Cruz P., Theau J.-P., Lecloux E., Jouany C., Duru M. (2010) : "Typologie fonctionnelle de graminées fourragères pérennes : une classification multitraits ", *Fourrages*, 201, 11-17.

Duru M., Justes E., Langlet A., Tirilly V., (1993) : "Comparaison des dynamiques d'apparition et de mortalité des organes de fétuque élevée, dactyle et luzerne (feuilles, talles et tiges)", *Agronomie*, 13, 237-252

Duru M., Calvière I., Balent G., Langlet A., (1993) : "Pédoclimat, fertilisation et croissance des prairies permanentes au printemps", *Fourrages*, 133, 43-57

Duru M., Balent G., Gibon A., Magda D., Theau JP., Cruz P., Jouany C. (1998) : "Fonctionnement et dynamique des prairies permanentes, Exemple des Pyrénées centrales", *Fourrages*, 153, 97-113.

Duru M., Hazard L., Jeangros B., Mosimann E. (2001) : "Fonctionnement de la prairie pâturée : structure du couvert et biodiversité", *Fourrages*, 166, 165-188

Galliot J.N., Hulin S., Le Henaff P.M., Farruggia A., Seytre L., Perera S., Dupic G., Faure P., Carrère P., 2020. "Typologie multifonctionnelle des prairies du Massif central", *Edition Sidam-AEOLE*, 284 pages.

Programme Herbe et Fourrages Centre : (2014) "Guide du pâturage"

Mariotte P et al., (2022) "Utiliser la somme des températures pour évaluer le rendement et la qualité fourragère des prairies permanentes et temporaires intensives", *Agroscope*

Moreau JC. (2014) Guide Méthodologique PraiCos « Optimiser le potentiel productif des prairies – Démarche de conseil n°4 élaborée dans le cadre du Casdar PraiCos » *IDELE*

Seuret J.-M., Theau J.-P., Pottier E., Pelletier P., Piquet M., Delaby L., (2014) : "Des outils d'aide à la gestion du pâturage pour mieux valoriser les prairies et renforcer la confiance des éleveurs", *Fourrages*, 218, 191-201

Theau J.-P., Zeorourou A., (2008) : "Herb'âge, une méthode de calcul des sommes de températures pour la gestion des prairies", *Les Cahiers d'Orphée*, 91-102